


Claims

1. A method of treating a susceptible cancer in a human or animal subject mammal, comprising administering to said subject an effective amount of a compound of formula (I):



or a salt or solvate thereof;

wherein X is N or CH;

Y is a group W(CH<sub>2</sub>), (CH<sub>2</sub>)W, or W, in which W is O, S(O)<sub>m</sub> wherein m is 0, 1 or 2, or NR<sup>a</sup> wherein R<sup>a</sup> is hydrogen or a C<sub>1-8</sub> alkyl group;

R<sup>1</sup> represents a 5- or 6-membered heterocyclic ring containing 1 to 4 heteroatoms selected from N, O or S(O)<sub>m</sub>, wherein m is as defined above, with the provisos that the ring does not have two adjacent O or S(O)<sub>m</sub> atoms and that where the ring has only N as heteroatom(s) the ring is C-linked to the quinazoline or quinoline ring, R<sup>1</sup> being optionally substituted by one or more R<sup>3</sup> groups;

each R<sup>3</sup> is independently selected from the group consisting of amino, hydrogen, halogen, hydroxy, nitro, carboxy, formyl, cyano, trifluoromethyl, trifluoromethoxy, carbamoyl, ureido, guanidino, C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkoxy, C<sub>3-8</sub> cycloalkoxyl, C<sub>4-8</sub> alkylcycloalkoxy, C<sub>1-8</sub> alkylcarbonyl, C<sub>1-8</sub> alkoxy carbonyl, N-C<sub>1-4</sub> alkylcarbamoyl, N,N-di-[C<sub>1-4</sub> alkyl]carbamoyl, hydroxyamino, C<sub>1-4</sub> alkoxyamino, C<sub>2-4</sub> alkanoyloxyamino, C<sub>1-4</sub> alkylamino, di[C<sub>1-4</sub> alkyl]amino, di-[C<sub>1-4</sub> alkyl]amino-C<sub>1-4</sub> alkylene-(C<sub>1-4</sub> alkyl)amino, C<sub>1-4</sub> alkylamino- C<sub>1-4</sub>

alkylene-(C<sub>1-4</sub> alkyl)amino, hydroxy-C<sub>1-4</sub> alkylene-(C<sub>1-4</sub> alkyl)amino, phenyl, phenoxy, 4-pyridon-1-yl, pyrrolidin-1-yl, imidazol-1-yl, piperidino, morpholino, thiomorpholino, thiomorpholino-1-oxide, thiomorpholino-1,1-dioxide, piperazin-1-yl, 4-C<sub>1-4</sub> alkylpiperazin-1-yl, dioxolanyl, C<sub>1-8</sub> alkylthio, arylthio, C<sub>1-4</sub> alkylsulphinyl, C<sub>1-4</sub> alkylsulphonyl, arylsulphonyl, arylsulphinyl, halogeno-C<sub>1-4</sub> alkyl, hydroxy-C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkanoyloxy-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy-C<sub>1-4</sub> alkyl, carboxy-C<sub>1-4</sub> alkyl, formyl-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxycarbonyl-C<sub>1-4</sub>-alkyl, carbamoyl-C<sub>1-4</sub> alkyl, N-C<sub>1-4</sub> alkylcarbamoyl-C<sub>1-4</sub>alkyl, N,N-di-[C<sub>1-4</sub> alkyl]carbamoyl-C<sub>1-4</sub>alkyl, amino-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkylamino-C<sub>1-4</sub> alkyl, di-[C<sub>1-4</sub> alkyl]amino-C<sub>1-4</sub> alkyl, phenyl-C<sub>1-4</sub> alkyl, 4-pyridon-1-yl-C<sub>1-4</sub> alkyl, pyrrolidin-1-yl-C<sub>1-4</sub> alkyl, imidazol-1-yl-C<sub>1-4</sub> alkyl, piperidino-C<sub>1-4</sub> alkyl, morpholino-C<sub>1-4</sub> alkyl, thiomorpholino-C<sub>1-4</sub> alkyl, thiomorpholino-1-oxide-C<sub>1-4</sub>alkyl, thiomorpholino-1,1-dioxide-C<sub>1-4</sub>alkyl, piperazin-1-yl-C<sub>1-4</sub>alkyl, 4-C<sub>1-4</sub> alkylpiperazin-1-yl-C<sub>1-4</sub> alkyl, hydroxy-C<sub>2-4</sub> alkoxy-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy-C<sub>2-4</sub> alkoxy-C<sub>1-4</sub> alkyl, hydroxy-C<sub>2-4</sub> alkylamino-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy-C<sub>2-4</sub> alkylamino-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkylthio-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkylsulphinyl-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkylsulphonyl-C<sub>1-4</sub> alkyl, hydroxy-C<sub>2-4</sub> alkylthio-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy-C<sub>2-4</sub> alkylthio-C<sub>1-4</sub> alkyl, phenoxy-C<sub>1-4</sub> alkyl, anilino-C<sub>1-4</sub> alkyl, phenylthio-C<sub>1-4</sub> alkyl, cyano-C<sub>1-4</sub> alkyl, halogeno-C<sub>2-4</sub> alkoxy, hydroxy-C<sub>2-4</sub> alkoxy, C<sub>2-4</sub> alkanoyloxy-C<sub>2-4</sub> alkoxy, C<sub>1-4</sub> alkoxy-C<sub>2-4</sub> alkoxy, carboxy-C<sub>1-4</sub> alkoxy, formyl-C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkoxycarbonyl-C<sub>1-4</sub> alkoxy, carbamoyl-C<sub>1-4</sub> alkoxy, N-C<sub>1-4</sub> alkylcarbamoyl-C<sub>1-4</sub> alkoxy, N,N-di-[C<sub>1-4</sub> alkyl]carbamoyl-C<sub>1-4</sub> alkoxy, amino-C<sub>2-4</sub> alkoxy, C<sub>1-4</sub> alkylamino-C<sub>2-4</sub> alkoxy, di-[C<sub>1-4</sub> alkyl]amino-C<sub>2-4</sub> alkoxy, di-[C<sub>1-4</sub> alkyl-C<sub>2-4</sub> alkoxy]amino-C<sub>2-4</sub> alkoxy, C<sub>2-4</sub> alkanoyloxy, hydroxy-C<sub>2-4</sub> alkanoyloxy, C<sub>1-4</sub>alkoxy-C<sub>2-4</sub> alkanoyloxy, phenyl-C<sub>1-4</sub> alkoxy, phenoxy-C<sub>2-4</sub> alkoxy, anilino-C<sub>2-4</sub> alkoxy, phenylthio-C<sub>2-4</sub> alkoxy, 4-pyridon-1-yl-C<sub>2-4</sub> alkoxy, piperidino-C<sub>2-4</sub> alkoxy, morpholino-C<sub>2-4</sub> alkoxy,

thiomorpholino-C<sub>2-4</sub> alkoxy, thiomorpholino-1-oxide-C<sub>2-4</sub> alkoxy,  
 thiomorpholino-1,1-dioxide-C<sub>2-4</sub> alkoxy, piperazin-1-yl-C<sub>2-4</sub> alkoxy, 4-C<sub>1-4</sub>  
 alkylpiperazin-1-yl-C<sub>2-4</sub> alkoxy, pyrrolidin-1-yl-C<sub>2-4</sub> alkoxy, imidazol-1-yl-C<sub>2-4</sub>  
 alkoxy, halogeno-C<sub>2-4</sub> alkylamino, hydroxy-C<sub>2-4</sub> alkylamino, C<sub>2-4</sub>  
 alkanoyloxy-C<sub>2-4</sub> alkylamino, C<sub>1-4</sub> alkoxy-C<sub>2-4</sub> alkylamino, carboxy-C<sub>1-4</sub>  
 alkylamino, C<sub>1-4</sub> alkoxycarbonyl-C<sub>1-4</sub> alkylamino, carbamoyl-C<sub>1-4</sub>  
 alkylamino, N-C<sub>1-4</sub> alkylcarbamoyl-C<sub>1-4</sub> alkylamino, N,N-di-[C<sub>1-4</sub>  
 alkyl]carbamoyl-C<sub>1-4</sub> alkylamino, amino-C<sub>2-4</sub> alkylamino, C<sub>1-4</sub> alkylamino-  
 C<sub>2-4</sub> alkylamino, di-[C<sub>1-4</sub>alkyl]amino-C<sub>2-4</sub> alkylamino, phenyl-C<sub>1-4</sub>  
 alkylamino, phenoxy-C<sub>2-4</sub> alkylamino, anilino-C<sub>2-4</sub> alkylamino, 4-pyridon-1-  
 yl- C<sub>2-4</sub> alkylamino, pyrrolidin-1-yl-C<sub>2-4</sub> alkylamino, imidazol-1-yl-C<sub>2-4</sub>  
 alkylamino, piperidino-C<sub>2-4</sub> alkylamino, morpholino-C<sub>2-4</sub> alkylamino,  
 thiomorpholino-C<sub>2-4</sub> alkylamino, thiomorpholino-1-oxide-C<sub>2-4</sub> alkylamino,  
 thiomorpholino-1,1-dioxide-C<sub>2-4</sub> alkylamino, piperazin-1-yl-C<sub>2-4</sub>alkylamino,  
 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl-C<sub>2-4</sub>alkylamino, phenylthio-C<sub>2-4</sub> alkylamino, C<sub>2-4</sub>  
 alkanoylamino, C<sub>1-4</sub> alkoxycarbonylamino, C<sub>1-4</sub> alkylsulphonylamino, C<sub>1-4</sub>  
 alkylsulphinylamino, benzamido, benzenesulphonamido, 3-phenylureido, 2-  
 oxopyrrolidin-1-yl, 2,5-dioxopyrrolidin-1-yl, halogeno-C<sub>2-4</sub> alkanoylamino,  
 hydroxy-C<sub>2-4</sub> alkanoylamino, hydroxy-C<sub>2-4</sub> alkanoyl-(C<sub>1-4</sub> alkyl)-amino, C<sub>1-4</sub>  
 alkoxy-C<sub>2-4</sub> alkanoylamino, carboxy-C<sub>2-4</sub> alkanoylamino, C<sub>1-4</sub>  
 alkoxycarbonyl-C<sub>2-4</sub> alkanoylamino, carbamoyl-C<sub>2-4</sub> alkanoylamino, N-C<sub>1-4</sub>  
 alkylcarbamoyl-C<sub>2-4</sub> alkanoylamino, N,N-di-[C<sub>1-4</sub> alkyl]carbamoyl-C<sub>2-4</sub>  
 alkanoylamino, amino-C<sub>2-4</sub> alkanoylamino, C<sub>1-4</sub> alkylamino-C<sub>2-4</sub>  
 alkanoylamino or di-[C<sub>1-4</sub> alkyl]amino-C<sub>2-4</sub> alkanoylamino; and wherein said  
 benzamido or benzenesulphonamido substituent or any anilino, phenoxy or  
 phenyl group on a R<sup>3</sup> substituent may optionally have one or two halogeno,  
 C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy substituents; and wherein any substituent having a  
 heterocyclic ring may optionally have one or two halogeno, C<sub>1-4</sub> alkyl or C<sub>1-4</sub>  
 alkoxy substituents on said ring; and wherein any substituent having a

Sub  
B2

This document contains information that is exempt from disclosure under the Freedom of Information Act (5 U.S.C. 552). This information is being furnished to you for your information only. It is not to be released to the public.

heterocyclic ring may optionally have one or two oxo or thioxo substituents on said ring;

or  $R^3$  represents a group selected from  $M^1-M^2-M^3-M^4$ ,  $M^1-M^5$  or  $M^1-M^2-M^3-M^6$  wherein

$M^1$  represents a  $C_{1-4}$  alkyl group, wherein optionally a  $CH_2$  group is replaced by a CO group;


$M^2$  represents  $NR^{12}$  or  $CR^{12}R^{13}$ , in which  $R^{12}$  and  $R^{13}$  each independently represent H or  $C_{1-4}$  alkyl;

$M^3$  represents a  $C_{1-4}$  alkyl group;

$M^3'$  represents a  $C_{1-4}$  alkyl group or is absent;

$M^4$  represents  $CN$ ,  $NR^{12}S(O)_mR^{13}$ ,  $S(O)_mNR^{14}R^{15}$ ,  $CONR^{14}R^{15}$ ,  $S(O)_mR^{13}$  or  $CO_2R^{13}$ , in which  $R^{12}$ ,  $R^{13}$  and  $m$  are as defined above and  $R^{14}$  and  $R^{15}$  each independently represent H or  $C_{1-4}$  alkyl, or  $R^{14}$  and  $R^{15}$  together with the nitrogen atom to which they are attached form a 5- or 6-membered ring optionally containing 1 or 2 additional heteroatoms selected from N, O or S( $O$ ) $_m$  in which ring any nitrogen atom present may optionally be substituted with a  $C_{1-4}$  alkyl group, and which ring may optionally have one or two oxo or thioxo substituents;

$M^5$  represents the group  $NR^{14}R^{15}$ , wherein  $R^{14}$  and  $R^{15}$  are as defined above, or  $M^5$  represents the group



in which  $t$  represents 2 to 4 and  $R^{16}$  represents OH,  $OC_{1-4}$  alkyl or  $NR^{14}R^{15}$ ; and

$M^6$  represents a  $C_{3-6}$  cycloalkyl group, the group  $NR^{14}R^{15}$ , wherein  $R^{14}$  and  $R^{15}$  are as defined above, or a 5- or 6-membered heterocyclic ring system containing 1 to 4 heteroatoms selected from N, O or S;

and  $p$  is 0 to 3; or when  $p$  is 2 or 3, two adjacent  $R^3$  groups together form an optionally substituted methylenedioxy or ethylenedioxy group;


*Sub B2*

$R^2$  is selected from the group consisting of hydrogen, halogen, trifluoromethyl,  $C_{1-4}$  alkyl and  $C_{1-4}$  alkoxy;

U represents phenyl or a 5 to 10-membered mono or bicyclic ring system in which one or more of the carbon atoms is optionally replaced by a heteroatom independently selected from N, O and  $S(O)_m$ , wherein m is 0, 1 or 2, and wherein U is substituted by at least one independently selected  $R^6$  group and U is optionally substituted by at least one independently selected  $R^4$  group;

each  $R^4$  is independently hydrogen, hydroxy, halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  alkylamino, di- $[C_{1-4}$  alkyl]amino,  $C_{1-4}$  alkylthio,  $C_{1-4}$  alkylsulphinyl,  $C_{1-4}$  alkylsulphonyl,  $C_{1-4}$  alkylcarbonyl,  $C_{1-4}$  alkylcarbamoyl, di- $[C_{1-4}$  alkyl] carbamoyl, carbamyl,  $C_{1-4}$  alkoxy carbonyl, cyano, nitro or trifluoromethyl;

each  $R^6$  is independently a group  $ZR^7$  wherein Z is joined to  $R^7$  through a  $(CH_2)_p$  group in which p is 0, 1 or 2 and Z represents a group  $V(CH_2)$ ,  $V(CF_2)$ ,  $(CH_2)V$ ,  $(CF_2)V$ ,  $V(CRR')$ ,  $V(CHR)$  or V where R and R' are each  $C_{1-4}$  alkyl and in which V is a hydrocarbonyl group containing 0, 1 or 2 carbon atoms, carbonyl, dicarbonyl,  $CH(OH)$ ,  $CH(CN)$ , sulphonamide, amide, O,  $S(O)_m$  or  $NR^b$  where  $R^b$  is hydrogen or  $R^b$  is  $C_{1-4}$  alkyl; and  $R^7$  is an optionally substituted  $C_{3-6}$  cycloalkyl; or an optionally substituted 5, 6, 7, 8, 9 or 10-membered carbocyclic or heterocyclic moiety; or  $R^6$  is a group  $ZR^7$  in which Z is  $NR^b$ , and  $NR^b$  and  $R^7$  together form an optionally substituted 5, 6, 7, 8, 9 or 10-membered carbocyclic or heterocyclic moiety.

2. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible breast cancer.
3. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible non-small cell lung cancer.
4. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible ovarian cancer.
5. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible stomach cancer.
6. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible pancreatic cancer.
7. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible head and neck cancer.
8. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible cancer characterized by expression or over-expression of EGFR.
9. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible cancer characterized by expression or over-expression of erbB-2.
10. A method as claimed in claim 1, wherein the susceptible cancer is a susceptible cancer characterized by expression 

Add  
B3